



South Dakota

Live births, abortions, and pregnancies


Live birth rates, 1982–1992



Live birth rates,* 1991



Pregnancy rates, 1990



National live birth rate ranking, 1990


Age group		
15-17	18-19	15-19
43rd highest	33rd highest	37th highest

*Missing bar indicates fewer than 10,000 females or fewer than 20 births. This graph refers to females aged 15–19 years.




Sexually transmitted diseases South Dakota

Gonorrhea rates,* 1982–1992



Gonorrhea rates, 1992



National ranking,[†] 1992

Males – 45th highest Females – 43rd highest


*Data were unavailable before 1986.

†National ranking refers to males and females aged 10–24 years.




Chlamydia cases and rates,* 1988–1992


Cases



Rates




Cases



Rates



Acquired immunodeficiency syndrome (AIDS) cumulative cases,† 1981–1992



*Chlamydia cases and rates refer to males and females of all ages.


†Missing bar indicates fewer than six cumulative cases.



Related risk behaviors,* 1993

South Dakota

Had sexual intercourse (lifetime)




Had sexual intercourse with four or more people (lifetime)




Had sexual intercourse during the past three months (currently sexually active)



Used a condom during last sexual intercourse (among currently sexually active)



Used birth control pills during last sexual intercourse (among currently sexually active)



Injected illegal drugs (lifetime)



*Related risk behaviors refer to high school students.

Sources: CDC—NCHS and NCCDPHP (birth, abortion, and pregnancy data), NCPS (gonorrhea and chlamydia data), NCID (AIDS data); 1993 South Dakota Youth Risk Behavior Survey (risk behavior data).

